

Table 27

Ranked Losses by pest

Cotton Insect losses 2010

pest	% Reduction	acres infested	rank	% infested	Cost/acre	Bales lost
Bollworm/Budworm	1.186%	8,148,844	1	76.44%	\$2.54	263,902
Stink Bugs	0.724%	6,712,988	2	62.97%	\$3.45	162,397
Lygus	0.677%	5,932,835	3	55.65%	\$6.86	191,826
Cotton Fleahopper	0.362%	4,487,032	4	42.09%	\$2.79	81,048
Aphids	0.286%	7,133,029	5	66.91%	\$1.39	60,377
Thrips	0.200%	10,165,601	6	95.36%	\$2.32	45,964
Spider Mites	0.199%	3,522,479	7	33.04%	\$1.75	57,189
Fall Armyworm	0.199%	2,762,701	8	25.92%	\$0.20	41,256
Clouded Plant bugs	0.024%	614,569	9	5.77%	\$0.14	6,465
Silverleaf Whitefly (Bemisia)	0.020%	508,430	10	4.77%	\$0.81	4,935
Other Insects*	0.024%	649,594	11	6.09%	\$0.02	5,076
Cutworms	0.003%	487,946	12	4.58%	\$0.19	699
Loopers	0.001%	735,998	13	6.90%	\$0.01	197
Grasshoppers	0.001%	1,293,128	14	12.13%	\$0.02	373
Beet Armyworm	0.001%	967,552	15	9.08%	\$0.06	146
Saltmarsh Caterpillars	0.001%	689,127	16	6.46%	\$0.00	140
Banded Winged Whitefly	0.000%	483,273	17	4.53%	\$0.00	0
Pink Bollworm	0.000%	97,725	18	0.92%	\$0.00	0
Cotton Leaf Perforator	0.000%	14,988	19	0.14%	\$0.00	0
Southern Armyworms	0.000%	242,500	20	2.27%	\$0.00	0
Boll Weevil	0.000%	115,470	21	1.08%	\$0.01	0
European Cornborer	0.000%	0	22	0.00%	\$0.00	0